Osteoporosis Management in Women Who Had a Fracture (OMW)

HEDIS® Measurement Year 2022 Measures

Measure Description: Assess the percentage of women 67-85 years of age who suffered a fracture between July 1 of the year prior to the measurement year, through June 30 of the measurement year, and who had either a bone mineral density test (BMD) or a prescription for a drug to treat osteoporosis in the six months (180 days) after the fracture.

Eligible Population

• Females aged 67-85 through December 31 of the measurement year (MY).

Exclusions

- BMD within 24 months prior to Index Episode Start Date (IESD)*.
- Osteoporosis therapy or dispensed prescription medication within 12 months prior to IESD.
- Hospice
- Fractures of finger, toe, face and skull.
- Members 66 years of age and older as of December 31 of the MY with frailty and advanced illness during the MY.

*IESD: the earliest date of service with a fracture diagnosis within the time frame specified above (July-June).

National Osteoporosis Foundation Resources

The National Osteoporosis Foundation's website, <u>NOF.Org</u>, offers a variety of resources and tools for health care professionals, including information about:

- Prevention of osteoporosis and broken bones.
- Promotion of strong bones for life.
- Education, advocacy, and research.

Numerator Codes

The complete NCQA approved code set list can be referenced in the coding guide at https://www.healthplan.org/providers/patientcare-programs/quality-measures. For questions, please contact your practice management consultant. To identify your practice management consultant please refer to healthplan.org/providers/overview/meetprovider-servicing-team

Code Type	Codes	Code Description
CPT	76977, 77078, 77080, 77081, 77085, 77086	BMD Tests
ICD-10	BP48ZZ1, BP49ZZ1, BP4GZZ1, BP4HZZ1, BP4LZZ1, BP4MZZ1, BP4NZZ1, BP4PZZ1, BQ00ZZ1, BQ01ZZ1, BQ03ZZ1, BQ04ZZ1, BR00ZZ1, BR07ZZ1, BR09ZZ1, BR0GZZ1	BMD Tests
ICD-9	88.98	BMD Tests
HCPCS	J0897, J1740, J3110, J3111, J3489	Osteoporosis Medication Therapy



Osteoporosis Management in Women Who Had a Fracture (OMW)

HEDIS® Measurement Year 2022 Measures

Tips for Correct Coding

- Differentiate between active fractures and after care treatment.
- Active fracture treatment is **not** usually provided in a primary care setting.
- If there is no evidence of an active fracture, the billing provider can submit a corrected claim to have the member removed from the measure.

The Health Plan has a team of member advocates, health coaches, social workers and nurses who can assist you and your patients to remove or overcomeany barriers to care through benefit assistance, community resource referrals or enrollment in a THP clinical program. To refer a patient who is a THP member for assistance, call **1.877.903.7504** and let us know what we can do to help your patient receive and adhere to your recommended plan of care.

